



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>




ELSEVIER

Contents lists available at ScienceDirect

Journal of Differential Equations

www.elsevier.com/locate/jde



Alexandroff–Bakelman–Pucci estimate and Harnack inequality for degenerate/singular fully non-linear elliptic equations

Cyril Imbert

Université Paris-Dauphine, CEREMADE, UMR CNRS 7534, place de Lattre de Tassigny, 75775 Paris cedex 16, France

ARTICLE INFO

Article history:

Received 28 May 2010

Available online 5 August 2010

MSC:

35B45

15B65

35J15

49L25

Keywords:

Degenerate fully non-linear elliptic equation

Singular fully non-linear elliptic equation

Non-divergence form

Alexandroff–Bakelman–Pucci estimate

Weak Harnack inequality

Local maximum principle

Harnack inequality

Hölder regularity

Viscosity solutions

ABSTRACT

In this paper, we study fully non-linear elliptic equations in non-divergence form which can be degenerate or singular when “the gradient is small”. Typical examples are either equations involving the m -Laplace operator or Bellman–Isaacs equations from stochastic control problems. We establish an Alexandroff–Bakelman–Pucci estimate and we prove a Harnack inequality for viscosity solutions of such non-linear elliptic equations.

© 2010 Elsevier Inc. All rights reserved.

1. Introduction

Following the original strategy of Krylov and Safonov [22,23], Delarue [12] proved by probabilistic methods a Harnack inequality for quasi-linear elliptic equations of the form

$$-\operatorname{Tr}(A(x, u, Du)D^2u) + H(x, u, Du) = 0, \quad x \in \Omega \quad (1)$$

E-mail address: imberty@ceremade.dauphine.fr.

(where Ω is a domain of \mathbb{R}^n) in the case where the $n \times n$ matrix $A(x, p)$ can degenerate. Precisely, he assumes

$$\Lambda^{-1}\lambda(p)I \leq A(x, u, p) \leq \Lambda\lambda(p)I, \tag{2}$$

$$H(x, u, p) \leq \Lambda(1 + \lambda(p))(1 + |p|) \tag{3}$$

where $\Lambda \geq 1$, $\lambda: \mathbb{R}^n \rightarrow \mathbb{R}^+$ is continuous and such that $\lambda(p) \geq \lambda_F$ if $|p| \geq M_F$. In (2), I denotes the identity matrix and the inequalities are understood in the sense of the usual partial order on the set of real symmetric matrices. The model example of (1) is the m -Laplace equation where $A(x, p) = |p|^{m-2}$ for some $m > 2$. An important application of the Harnack inequality is the derivation of a Hölder estimate for the solution of (1).

In this paper, we generalize this result to the case of fully non-linear elliptic equations in non-divergence form

$$F(x, u, Du, D^2u) = 0, \quad x \in \Omega, \tag{4}$$

which can be either degenerate or singular. We do so by proving first an Alexandroff–Bakelman–Pucci (ABP for short) estimate. This is the first main difference with [12] and the first main contribution of this paper. Important examples of (4) which are out of the scope of [12] are Bellman–Isaacs equations appearing in the context of stochastic control problems. We also generalize and/or recover results from [10,3] where an ABP estimate and a Harnack inequality respectively are obtained for

$$F_0(Du, D^2u) + b(x) \cdot Du|Du|^\alpha + cu|u|^\alpha + f_0(x) = 0, \quad x \in \Omega, \tag{5}$$

where F is positively homogeneous of order $\alpha \in (-1, 1)$ (see Section 6 for precise assumptions). If $\alpha \in [0, 1)$, the equation is degenerate. If $\alpha \in (-1, 0]$, the equation is singular. Even if this equation does not formally enter into our general framework, we will explain how the results of [10,3] can be derived from ours.

Known results. Krylov and Safonov [22,23] first proved a Harnack inequality for second order elliptic equations in non-divergence form with measurable coefficients. This result is often presented as the counterpart of the De Giorgi and Nash estimates [11,25] for divergence form equations.

As far as degenerate elliptic equations are concerned, De Giorgi and Nash estimates were obtained for equations in divergence form and for degeneracies of p -Laplace type. See for instance [27,24].

Krylov and Safonov estimates were obtained by Caffarelli [5] for fully non-linear elliptic equations of the form $F(x, D^2u) = 0$ (see also [29,16]). As explained in [6], a fundamental tool in this approach is the Alexandroff–Bakelman–Pucci estimate. Many authors extended these results since then; see for instance [14,20,7,26] and references therein.

To the best of our knowledge and as far as degenerate elliptic equations in non-divergence form are concerned, the Krylov and Safonov estimates obtained by Delarue [12] are the first ones.

After this work was completed, Birindelli and Demengel [3] obtained a Harnack inequality for degenerate elliptic equations of the form (5) with $\alpha \in [0, 1)$ in dimension 2. Reading their interesting paper, we understood that we could recover (and in fact extend) their results and deal with singular equations. We will explain how to get the same estimate in any dimension (see Section 6). Their work aims at generalizing the results of Dávila, Felmer and Quaas [9] where the same elliptic equation is considered but with $\alpha \in (-1, 0]$. Hence, the equation is singular. We also mention that an ABP estimate is proved in [10] for degenerate and singular equations. We will explain that it can be derived from ours; see Section 6 where our results are compared with the ones in [3,10].

Main results. Let us now describe a bit more precisely our main results. We use the techniques developed by Caffarelli [5] (see also [6]) instead of probability arguments to get, apart from the Alexandroff–Bakelman–Pucci estimate, a weak Harnack inequality and a local maximum principle. It is then easy to derive a Harnack inequality and a Hölder estimate of a solution of (4).

First and foremost, we mention that, as in [5,12], we use the notion of viscosity solution [8] since the equation is fully non-linear. We recall that if singular equations of the form (5) are considered, the classical notion of viscosity solutions must be adapted; see [2].

We next make precise the standing assumptions that the non-linearity F must satisfy. Throughout the paper, \mathcal{S}_n denotes the space of real symmetric $n \times n$ matrices and B_R denotes the open ball of radius $R \geq 0$.

Assumption (A).

- F is continuous on $\Omega \times \mathbb{R} \times \mathbb{R}^n \setminus B_{M_F} \times \mathcal{S}_n$ for some $M_F \geq 0$;
- F is (degenerate) elliptic, i.e. for all $x \in \Omega$, $r \in \mathbb{R}$, $p \in \mathbb{R}^n$ ($p \neq 0$ for singular equation) and $X, Y \in \mathcal{S}_n$,

$$X \leq Y \quad \Rightarrow \quad F(x, r, p, Y) \leq F(x, r, p, X).$$

- F is proper, i.e. it is non-decreasing with respect to its r variable.

Our first main result (Theorem 1) is an ABP estimate for lower semi-continuous super-solutions of (4) on a ball B_d where F is strictly elliptic for “large gradients”

$$\left. \begin{array}{l} X \geq 0 \\ |p| \geq M_F \\ F(x, r, p, X) \geq 0 \end{array} \right\} \Rightarrow -\lambda_F \operatorname{tr}(X) + \sigma(x)|p| + g(x, u) \geq 0 \tag{6}$$

for some continuous functions g and σ and some constants $M_F \geq 0$, $\lambda_F > 0$. This condition holds true if F satisfies (7) but it is more general. An ABP estimate permits us to control $\sup_{B_d} u^-$ in terms of $M_\partial = \sup_{\partial B_d} u^-$ and the L^n -norms of $g(x, M_\partial)$ and σ appearing in (6). In order to get such an estimate, we use the techniques from [5]. As we already mentioned it in [17], the ABP estimate that we are able to obtain differs slightly from classical ones in the sense that we can prove it under a weaker condition than (7); moreover, the super-solution is only lower semi-continuous. We recall that this is an a priori estimate: structure conditions ensuring the uniqueness of the solution are not required. We finally mention that when the equation is strictly elliptic ($M_F = 0$), we recover the classical ABP estimate.

Our second main result (Corollary 1) is a Harnack inequality for (4). This inequality is a consequence of a weak Harnack inequality and a local maximum principle proved by generalizing in an appropriate way (2) and (3). In view of (2), one can consider the quasi-linear equation (1) where A and H are replaced with

$$\tilde{A}(x, u, Du) = \frac{1}{\lambda(Du)} A(x, u, Du) \quad \text{and} \quad \tilde{H}(x, u, Du) = \frac{1}{\lambda(Du)} H(x, u, Du).$$

Hence, the new quasi-linear equation is uniformly elliptic. However, the first order term is, in this case, eventually singular and (2) can be seen as an assumption concerning the first order term. In the case of the m -Laplace equation, $\lambda(p) = |z|^{m-2}$ and H has therefore a polynomial growth of order $m - 1$. Assumptions (2), (3) are replaced with

$$\left. \begin{array}{l} |p| \geq M_F \\ F(x, u, p, X) \geq 0 \end{array} \right\} \Rightarrow \mathcal{M}^+(X) + \sigma(x)|p| + \gamma_F u + f(x) \geq 0, \tag{7}$$

$$\left. \begin{array}{l} |p| \geq M_F \\ F(x, u, p, X) \leq 0 \end{array} \right\} \Rightarrow \mathcal{M}^-(X) - \sigma(x)|p| + \gamma_F u - f(x) \leq 0 \tag{8}$$

where $\sigma, f : \bar{B} \rightarrow \mathbb{R}$ are continuous and M_F and γ_F are non-negative constants. It is important to remark that if F satisfies (7), (8), then it can be degenerate or singular and it can have a superlinear growth in p .

An important consequence of the Harnack inequality is the Hölder regularity of solutions of (4) (see Theorem 2). As far as the regularity of solutions of (4) is concerned, we notice that by assuming (7) and (8), we cannot expect more than Lipschitz continuity. Indeed, by making such an assumption, we somehow forget about all small gradients and we cannot expect these small gradients to be regular. We also point out that it is easier to prove the uniqueness of a Hölder continuous function than to prove a strong comparison result between discontinuous viscosity sub- and super-solutions (which is the classical way to get uniqueness of viscosity solutions [8]). To finish with, we shed light on the fact that, as for the ABP estimate, we recover the Harnack inequality of [5] in the strictly elliptic case ($M_F = 0$).

Extensions. We will explain how to deal with non-linearities, after redefining them if necessary, growing quadratically with respect to the gradient. Precisely, (7) and (8) are replaced with

$$\left. \begin{array}{l} |p| \geq M_F \\ F(x, u, p, X) \geq 0 \end{array} \right\} \Rightarrow \mathcal{M}^+(X) + \sigma(x)|p| + \sigma_2|p|^2 + \gamma_F u + f(x) \geq 0, \quad (9)$$

$$\left. \begin{array}{l} |p| \geq M_F \\ F(x, u, p, X) \leq 0 \end{array} \right\} \Rightarrow \mathcal{M}^-(X) - \sigma(x)|p| - \sigma_2|p|^2 + \gamma_F u - f(x) \leq 0 \quad (10)$$

where $\sigma, f : \bar{B} \rightarrow \mathbb{R}$ are continuous and M_F, σ_2 and γ_F are non-negative constants. In this case, it is known [29,21] that it is not possible to get a weak Harnack inequality which does not depend on the L^∞ -norm of the solution. See Section 5 for more details and comments.

As far as extensions of these results are concerned, we would like to mention next that we could have used L^p -viscosity solutions [4] instead of classical continuous viscosity solutions in order to be able to deal with discontinuous coefficients. We chose not to do so in order to avoid technicalities but we think that this can be done. We also mention that it is sometimes more difficult to get a classical ABP estimate when using this notion of solution; for instance in [20], the ABP estimate does not involve the contact set of the function.

We also mention that the parabolic case will be addressed in a future work.

Additional comments. Assumption (6) permits to take into account non-linearity growing linearly with respect to the gradient. Such an assumption appears in [28] where Trudinger proved that strong solutions satisfy a weak Harnack inequality for such non-linearities if σ is sufficiently integrable. This result has been generalized to viscosity solutions since then; see for instance [15,19].

We recall that it is possible to use the techniques introduced in [18] in order to prove the Hölder regularity of viscosity solutions much more easily. But the estimate of the Hölder constant depends in this case on the modulus of continuity of the coefficients of the equation.

Organization of the article. The paper is organized as follows. In Section 2, we construct a barrier function that will be used when proving the Harnack inequality. We also recall the definition of two Pucci operators. In Section 3, we establish an ABP estimate. In Section 4, we successively prove a weak Harnack inequality and a local maximum principle. We also derive from these two results a Harnack inequality. In Section 5, we explain how to deal with elliptic equations with quadratic dependence on the gradient. As applications of our results, we generalize and/or recover some results from [3,10] in Section 6. Section 7 is dedicated to proofs of our main results. Appendix A is added for the sake of completeness of proofs and for the reader's convenience. In particular, we give in Appendix A detailed proofs of results which can be easily derived from classical ones.

Notation. A ball of radius r centered at x is denoted by $B(x, r)$ or $B_r(x)$. If $x = 0$, we simply write B_r . ω_n denotes the volume of the unit ball. The hypercube $\prod_{i=1}^n (x_i - r/2, x_i + r/2)$ is denoted by $Q_r(x)$. If $x = 0$, we simply write Q_r .

Given a vector $a \neq 0$, \hat{a} denotes $a/|a|$. I denotes the identity matrix. The set of real symmetric $n \times n$ matrices is denoted by S_n .

A constant is universal if it only depends on n (dimension), q (constant greater than n fixed in all the paper), λ_F and Λ_F (ellipticity constants).

Given a lower semi-continuous function u , $D^{2,-}u(x)$ (resp. $D^{2,+}u(x)$) denotes the set of all subsets (resp. limiting subsets) of u at point x . See [8] for definitions.

2. Preliminaries

Pucci operators. We recall the definition of two important second order non-linear elliptic operators. For all $M \in S_n$, we define

$$\mathcal{M}^+(M) = \sup_{A \in \mathcal{A}_{\lambda_F, \Lambda_F}} (-\text{Tr}(AM)),$$

$$\mathcal{M}^-(M) = \inf_{A \in \mathcal{A}_{\lambda_F, \Lambda_F}} (-\text{Tr}(AM))$$

where $\mathcal{A}_{\lambda_F, \Lambda_F} = \{A \in S_n: \lambda_F I \leq A \leq \Lambda_F I\}$. We will refer to these operators as the *maximal* and *minimal Pucci operators*. Remark that \mathcal{M}^+ is subadditive. More precisely, it is the support function of the set $-\mathcal{A}_{\lambda_F, \Lambda_F}$. We will also use the fact that $\mathcal{M}^-(M) = -\mathcal{M}^+(-M)$.

Construction of a barrier. We now construct a barrier that will be used when proving the (weak) Harnack inequality.

Lemma 1 (*Construction of a barrier*). *Given a constant $\varepsilon_0 > 0$, there exist a smooth function $\varphi: \mathbb{R}^n \rightarrow \mathbb{R}$, a universal constant $M_B > 1$ and constants $C_B > 0$, $R, r > 0$ (with $R \geq (3r/2)\sqrt{n}$) depending only on the dimension n , λ_F , Λ_F and ε_0 , such that*

$$\varphi \geq 0 \quad \text{in } \mathbb{R}^n \setminus B_R, \tag{11}$$

$$\varphi \leq -2 \quad \text{in } Q_{3r}, \tag{12}$$

$$\varphi \geq -M_B \quad \text{in } \mathbb{R}^n, \tag{13}$$

$$|D\varphi| \leq \varepsilon_0 \quad \text{in } \mathbb{R}^n, \tag{14}$$

$$\mathcal{M}^-\varphi + C_B \xi \geq 0 \quad \text{in } \mathbb{R}^n \tag{15}$$

where $\xi: \mathbb{R}^n \rightarrow [0, 1]$ is a continuous function supported in \overline{Q}_r .

Remark 1. We recall that this barrier function will be used to prove the weak Harnack inequality. At first glance, it is not clear why we need to construct a function φ such that $\mathcal{M}^-\varphi \geq 0$ on Q_r and $\varphi \leq -2$ on Q_{3r} . This will be clearer when applying the cube decomposition in order to estimate the volume of all the level sets (and not only one) of a super-solution. And we choose $R \geq (3r/2)\sqrt{n}$ in order that $Q_{3r} \subset B_R$.

Proof of Lemma 1. We follow [6] by choosing φ under the following form for $x \notin B_r$

$$\varphi(x) = M_1 - M_2|x|^{-\alpha}$$

where $\alpha > 0$ will be chosen later and $M_1, M_2 > 0$ have to be chosen such that (11), (12) and (14) hold true for $x \notin B_r$ (with $R \geq (3r/2)\sqrt{n}$). It is enough to impose

$$\begin{aligned} M_2 &\leq M_1 R^\alpha, \\ ((3r/2)\sqrt{n})^\alpha (M_1 + 2) &\leq M_2, \\ M_2 &\leq \varepsilon_0 \frac{r^{\alpha+1}}{\alpha} \end{aligned}$$

or equivalently

$$((3r/2)\sqrt{n})^\alpha (M_1 + 2) \leq M_2 \leq \min(M_1 R^\alpha, \varepsilon_0 r^{\alpha+1}/\alpha).$$

One can choose M_2 and M_1 so that they satisfy the previous condition if and only if

$$2 \frac{((3r/2)\sqrt{n})^\alpha}{R^\alpha - ((3r/2)\sqrt{n})^\alpha} \leq M_1 \leq \frac{\varepsilon_0}{\alpha((3/2)\sqrt{n})^\alpha} r - 2.$$

Hence, we choose $R = q(3r/2)\sqrt{n}$ with $q > 1$ and $r > 0$ satisfying

$$\frac{2}{q^\alpha - 1} \leq \frac{\varepsilon_0}{\alpha((3/2)\sqrt{n})^\alpha} r - 2.$$

It is now enough to choose $q > 1$ such that $\frac{2}{q^\alpha - 1} \leq 1$ and r such that

$$\frac{\varepsilon_0}{\alpha((3/2)\sqrt{n})^\alpha} r \geq 3.$$

We now choose $\alpha > 0$ so that (15) holds true. If $x \notin B_r$, we have

$$\mathcal{M}^-(D^2\varphi(x)) = -\alpha M_2 |x|^{-(\alpha+2)} (\Lambda_F(n-1) - \lambda_F(\alpha+1)).$$

Hence it is enough to choose $\alpha > \max(0, \frac{\Lambda_F}{\lambda_F}(n-1) - 1)$ to conclude.

It is next easy to extend φ on \mathbb{R}^n such that (12) and (14) remain true and (13) is satisfied too for some universal constant $M_B > 1$. Indeed, we have outside B_r

$$\varphi \geq M_1 - M_2 r^{-\alpha} \geq 2 \frac{1}{q^\alpha - 1} - \frac{\varepsilon_0 r}{\alpha}.$$

It is now enough to remark that q and $\varepsilon_0 r$ can be chosen universal and we also saw above that α can be chosen universal too. Hence M_B can be chosen universal. \square

Rescaling solutions. We will have to rescale sub- or super-solutions several times. We need to know how non-linearities are rescaled in order, for instance, to determine if these new F 's satisfy assumptions.

Lemma 2 (*Rescaling solutions*). Given $R_0 > 0$, $t_0 > 0$ and $x_0 \in \mathbb{R}^n$, let u be a super-solution of F on $Q_{t_0 R_0}(x_0)$. Consider the linear map $T : Q_{R_0} \rightarrow Q_{t_0 R_0}(x_0)$ defined by $T(y) = x_0 + t_0 y$. Then the scaled solution $u_s(y) = \frac{1}{M_0} u(T(y))$ is a super-solution of $F_s = 0$ in Q_{R_0} with

$$F_s(y, v, q, Y) = \frac{t_0^2}{M_0} F(x_0 + t_0 y, M_0 v, t_0^{-1} M_0 q, t_0^{-2} M_0 Y).$$

If F satisfies (7) (resp. (8)), then F_s satisfies (7) (resp. (8)) with constants M_s, γ_s and functions σ_s and f_s

$$M_s = \frac{t_0 M_F}{M_0}, \quad \gamma_s = t_0^2 \gamma_F, \quad \sigma_s = t_0 \cdot \sigma \circ T, \quad f_s = \frac{t_0^2}{M_0} f \circ T.$$

In particular,

$$\|f_s\|_{L^n(Q_{R_0})} = \frac{t_0}{M_0} \|f\|_{L^n(Q_{t_0 R_0}(x_0))}, \quad \|\sigma_s\|_{L^q(Q_{R_0})} = t_0^{1-\frac{n}{q}} \|\sigma\|_{L^q(Q_{t_0 R_0}(x_0))}.$$

3. An ABP estimate

As explained in the Introduction, we can prove an ABP estimate as soon as the non-linearity F satisfies a strict ellipticity condition “for large gradients”. We must also prescribe a growth condition with respect to first order terms. We thus assume that F satisfies (6). Our first main result is the following theorem.

Theorem 1 (ABP estimate). Consider a non-linearity F which satisfies (A) and (6). Let u be a (lsc) super-solution of (4) in B_d . Then

$$\sup_{B_d} u^- \leq \sup_{\partial B_d} u^- + Cd \left(M_F + \left(\int_{B_d \cap \{u + M_\partial = \Gamma(u)\}} (f^+)^n \right)^{1/n} \right) \tag{16}$$

where $M_\partial = \sup_{\partial B_d} u^-$, $\Gamma(u)$ is the convex hull of $\min(u + M_\partial, 0)$ extended by 0 on B_{2d} , $f(x) = g(x, -M_\partial)$ and C is a constant (only) depending on $\|\sigma\|_{L^n(B_d)}$, n and λ_F .

Remark 2. Remark that when the equation is not degenerate ($M_F = 0$), Eq. (16) corresponds to the classical ABP estimate.

Remark 3. The constant C equals $3e^{C_{ABP}(1+\|\sigma\|_{L^n(B_d)}^n)}$ where $C_{ABP} = \frac{2^{n-2}}{\omega_n \lambda_F^n} n$.

Sketch of the proof of Theorem 1. The proof follows the ideas of [6,17]. The key lemma is the following one.

Lemma 3. The function $\Gamma(u)$ is $C^{1,1}$ on $\mathcal{B} = \{x \in B_d : |D\Gamma(u)(x)| \geq M_F\}$.

Remark 4. Remark that before knowing that $\Gamma(u)$ is $C^{1,1}$, $D\Gamma(u)$ is not uniquely determined. Hence \mathcal{B} should be first defined as follows

$$\mathcal{B} = \{x \in B_d : \forall (p, A) \in D^{2,-} \Gamma(u)(x), |p| \geq M_F\}.$$

Lemma 3 is proved together with

Lemma 4. The Hessian of $\Gamma(u)$ satisfies on \mathcal{B} the following properties

1. $D^2\Gamma(u) = 0$ a.e. in $\mathcal{B} \setminus \{u + M_\partial = \Gamma(u)\}$;
2. $D^2\Gamma(u)(x) \leq \lambda_F^{-1} \{\sigma(x) |D\Gamma(u)(x)| + f^+(x)\} I$ a.e. in $\mathcal{B} \cap \{u + M_\partial = \Gamma(u)\}$.

Proofs of these two lemmata can be adapted from the classical ones by remarking that points x_i called by $x \in \mathcal{B}$ when computing the convex hull $\Gamma(u)$ (see Proposition 1 in Appendix A) satisfy $D\Gamma(u)(x_i) = D\Gamma(u)(x)$. In particular, $x_i \in \mathcal{B}$, i.e. $|D\Gamma(u)(x_i)| \geq M_F$ and consequently (6) can be used. The reader is referred to Appendix A where detailed proofs are given for his convenience.

Lemma 5. *The following inclusion holds true*

$$B_{M/(3d)}(0) \setminus B_{M_F}(0) \subset D\Gamma(u)(\mathcal{B}) \tag{17}$$

where M denotes $(\sup_{B_d} u^- - \sup_{\partial B_d} u^-)^+$ and $\mathcal{B} = \{x \in B_d : |D\Gamma(u)(x)| \geq M_F\}$.

Proof. This lemma is a consequence of the classical fact

$$B_{M/(3d)}(0) \subset D\Gamma(u)(B_d). \quad \square$$

From now on, we assume without loss of generality that $M/(3d) \geq M_F$. We then use Lemma 3 in order to apply the area formula (see [13, Theorem 3.2.5] and Remark 6 below) to the Lipschitz map $D\Gamma(u) : \mathcal{B} \rightarrow \mathbb{R}^n$ and to the function $g(p) = (|p|^{n/(n-1)} + \mu^{n/(n-1)})^{(1-n)}$ for some positive real number μ to be fixed later,

$$\int_{D\Gamma(u)(\mathcal{B})} g(p) dp = \int_{\mathcal{B}} g(D\Gamma(u)) \det D^2\Gamma(u).$$

On one hand, we can use Lemmata 4 and 5 in order to get

$$\begin{aligned} \int_{B_{M/(3d)}(0) \setminus B_{M_F}(0)} g(p) dp &\leq \int_{D\Gamma(u)(\mathcal{B})} g(p) dp \\ &\leq \int_{\mathcal{B}} g(D\Gamma(u)) \det D^2\Gamma(u) \\ &\leq \frac{1}{\lambda_F^n} \int_{\mathcal{B} \cap \{u+M_\partial=\Gamma(u)\}} g(D\Gamma(u)) (\sigma |D\Gamma(u)| + f^+)^n \\ &\leq \frac{1}{\lambda_F^n} \int_{\mathcal{B} \cap \{u+M_\partial=\Gamma(u)\}} (|\sigma|^n + \mu^{-n} (f^+)^n). \end{aligned}$$

If now one chooses μ such that $\mu^n = \int_{\mathcal{B} \cap \{u+M_\partial=\Gamma(u)\}} (f^+)^n$, we obtain from the inequality $g(p) \geq 2^{2-n}(|p|^n + \mu^n)^{-1}$ the following estimate

$$\begin{aligned} \frac{2^{2-n}}{n} \omega_n \ln \frac{(M/(3d))^n + \mu^n}{(M_F)^n + \mu^n} &= 2^{2-n} \omega_n \int_{M_F}^{M/d} \frac{r^{n-1} dr}{r^n + \mu^n} \\ &\leq \int_{B_{M/d}(0) \setminus B_{M_F}(0)} g(p) dp \\ &\leq \lambda_F^{-n} (1 + \|\sigma\|_n^n) \end{aligned}$$

where ω_n denotes the volume of the unit ball. It is now easy to get (16). \square

Remark 5. We see from the previous proof that Assumptions (A) and (6) on F are important in order to get the following property

$$\forall (p, A) \in D^{2,-}u(x): \left. \begin{array}{l} A \geq 0, u(x) \leq 0 \\ |p| \geq M_F \end{array} \right\} \Rightarrow \lambda_F \operatorname{Tr} A \leq \sigma(x)|p| + f(x). \quad (18)$$

As a matter of fact, the previous piece of information is the relevant one in order to get (16). Indeed, in Lemma 4, the second estimate can be rewritten as follows

$$\lambda_F D^2 \Gamma(u)(x) \leq \{ \sigma(x) |D\Gamma(u)| + f(x) \} I.$$

Remark 6. The area formula in [13] is stated for maps $G: \mathbb{R}^n \rightarrow \mathbb{R}^n$ that are Lipschitz continuous on \mathbb{R}^n (in our case). However, the result still holds true if G is only Lipschitz continuous on \mathcal{B} since it is always possible to extend it in a Lipschitz map \tilde{G} on \mathbb{R}^n with $G = \tilde{G}$ on \mathcal{B} .

4. Harnack inequality

In this section, we explain how to derive a Harnack inequality from the ABP estimate. As usual, we obtain it by deriving on one hand a weak Harnack inequality and on the other hand a local maximum principle for the fully non-linear equation (4).

In order to get a weak Harnack inequality and a local maximum principle respectively, condition (6) is strengthened by assuming (7) and (8) respectively.

The Harnack inequality is obtained as a combination of the weak Harnack inequality and the local maximum principle. Here are precise statements.

Theorem 2 (Weak Harnack inequality). Given $q > n$ and a non-linearity F satisfying (A) and (7) for some continuous functions f and σ in Q_1 , consider a non-negative super-solution u of (4) in Q_1 . Then

$$\|u\|_{L^{p_0}(Q_{1/4})} \leq C \left(\inf_{Q_{1/2}} u + \max(M_F, \|f\|_{L^n(Q_1)}) \right) \quad (19)$$

where $p_0 > 0$ is universal and C (only) depends on $n, q, \lambda_F, \Lambda_F, \gamma_F$ and $\|\sigma\|_{L^q(Q_1)}$.

Theorem 3 (Local maximum principle). Given $q > n$ and a non-linearity F satisfying (A) and (8) for some continuous functions f and σ on Q_1 , consider a sub-solution u of (4) in Q_1 . Then for any $p > 0$,

$$\sup_{Q_{1/4}} u \leq C(p) \left(\|u^+\|_{L^p(Q_{1/2})} + \max(M_F, \|f\|_{L^n(Q_1)}) \right) \quad (20)$$

where $C(p)$ is a constant (only) depending on $n, q, \lambda_F, \Lambda_F, \gamma_F, \|\sigma\|_{L^q(Q_1)}$ and p .

Combining these two results, we obtain the second main result of this paper.

Corollary 1 (Harnack inequality). Given $q > n$ and a non-linearity F satisfying (A), (7) and (8) for some continuous functions f and σ on Q_1 , consider a non-negative solution u of (4) in Q_1 . Then

$$\sup_{Q_{1/2}} u \leq C \left(\inf_{Q_{1/2}} u + \max(M_F, \|f\|_{L^n(Q_1)}) \right) \quad (21)$$

where C is a constant (only) depending on $n, q, \lambda_F, \Lambda_F, \gamma_F$ and $\|\sigma\|_{L^q(Q_1)}$.

An important consequence of Corollary 1 is the following regularity result.

Corollary 2 (Interior Hölder regularity). Given $q > n$ and a non-linearity F satisfying (A), (7) and (8) for some continuous functions f and σ on Q_1 , consider a solution u of (4) in Q_1 . Then u is α -Hölder continuous on $Q_{\frac{1}{2}}$ and

$$\sup_{\substack{x,y \in Q_{\frac{1}{2}} \\ x \neq y}} \frac{|u(x) - u(y)|}{|x - y|^\alpha} \leq C_\alpha (\|u\|_{L^\infty(Q_1)} + \max(M_F, \|f\|_{L^n(Q_1)} + \gamma_F \|u\|_{L^\infty(Q_1)})) \quad (22)$$

where α and C_α depend (only) on $n, q, \lambda_F, \Lambda_F, \gamma_F$ and $\|\sigma\|_{L^q(Q_1)}$.

5. Quadratic growth in Du

In this section, we extend the results of the previous section to elliptic equations with a first order term (after changing the original equation if necessary; see the Introduction) which can grow quadratically with respect to the gradient. Precisely, (7) and (8) are replaced with (9) and (10).

Through a Cole–Hopf transform, an immediate consequence of Theorems 2 and 3 are the following results.

Theorem 4 (Weak Harnack inequality). Given $q > n$ and a non-linearity F satisfying (A) and (9) for some continuous functions f and σ in Q_1 , consider a non-negative super-solution u of (4) in Q_1 . Then

$$\|u\|_{L^{p_0}(Q_{1/4})} \leq C \left(\inf_{Q_{1/2}} u + \max(M_F, \|f\|_{L^n(Q_1)}) \right) \quad (23)$$

where $p_0 > 0$ is universal and C (only) depends on $\|u\|_{L^\infty(Q_1)}, n, q, \lambda_F, \Lambda_F, \gamma_F$ and $\|\sigma\|_{L^q(Q_1)}$.

Remark 7. As explained in [29,21], one cannot expect to get weak Harnack inequality for such non-linearities with a constant $C > 0$ which does not depend on a bound on u .

Remark 8. The constant C can be written

$$C = C_0 \frac{\frac{\sigma_2 \|u\|_{L^\infty(Q_1)}}{\lambda_F}}{1 - e^{-\frac{\sigma_2 \|u\|_{L^\infty(Q_1)}}{\lambda_F}}}$$

where C_0 (only) depends on $n, q, \lambda_F, \Lambda_F, \gamma_F$ and $\|\sigma\|_{L^q(Q_1)}$.

Theorem 5 (Local maximum principle). Given $q > n$ and a non-linearity F satisfying (A) and (10) for some continuous functions f and σ on Q_1 , consider a sub-solution u of (4) in Q_1 . Then for any $p > 0$,

$$\sup_{Q_{1/4}} u \leq C (\|u^+\|_{L^p(Q_{1/2})} + \max(M_F, \|f\|_{L^n(Q_1)})) \quad (24)$$

where C (only) depends on $\|u\|_{L^\infty(Q_1)}, n, q, \lambda_F, \Lambda_F, \gamma_F, \|\sigma\|_{L^q(Q_1)}$ and p .

Remark 9. The constant C can be written

$$C = C_0 \frac{\frac{\sigma_2 \|u\|_{L^\infty(Q_1)}}{\lambda_F}}{1 - e^{-\frac{\sigma_2 \|u\|_{L^\infty(Q_1)}}{\lambda_F}}}$$

where C_0 (only) depends on $n, q, \lambda_F, \Lambda_F, \gamma_F, \|\sigma\|_{L^q(Q_1)}$ and p .

It is now easy to derive a Harnack inequality and an interior Hölder estimate.

Corollary 3 (Harnack inequality). Given $q > n$ and a non-linearity F satisfying (A), (9) and (10) for some continuous functions f and σ on Q_1 , consider a non-negative solution u of (4) in Q_1 . Then

$$\sup_{Q_{1/2}} u \leq C \left(\inf_{Q_{1/2}} u + \max(M_F, \|f\|_{L^n(Q_1)}) \right) \tag{25}$$

where C (only) depends on $\|u\|_{L^\infty(Q_1)}$, n , q , λ_F , Λ_F , γ_F and $\|\sigma\|_{L^q(Q_1)}$.

Corollary 4 (Interior Hölder regularity). Given $q > n$ and a non-linearity F satisfying (A), (9) and (10) for some continuous functions f and σ on Q_1 , consider a solution u of (4) in Q_1 . Then u is α -Hölder continuous on $Q_{\frac{1}{2}}$ and

$$\sup_{\substack{x, y \in Q_{\frac{1}{2}} \\ x \neq y}} \frac{|u(x) - u(y)|}{|x - y|^\alpha} \leq C_\alpha \left(\|u\|_{L^\infty(Q_1)} + \max(M_F, \|f\|_{L^n(Q_1)} + \gamma_F \|u\|_{L^\infty(Q_1)}) \right) \tag{26}$$

where α and C_α depend (only) on n , q , λ_F , Λ_F , γ_F and $\|\sigma\|_{L^q(Q_1)}$.

6. Applications: results of [3,10]

In [3,10], Eq. (5) is considered. In [3], α lies in $[0, 1)$ and in [10], $\alpha > -1$. They assume

Assumption (H).

- (H1) $F_0(tp, \mu X) = |t|^\alpha \mu F_0(p, X)$ for $t \neq 0$ and $\mu \geq 0$ for some $\alpha > -1$;
- (H2) $|p|^\alpha \mathcal{M}^-(N) \leq F_0(p, M + N) - F_0(p, M) \leq |p|^\alpha \mathcal{M}^+(N)$.

The ABP estimate obtained in [10] is the following one

Theorem 6. (See [10, Theorem 1].) Under Assumption (H) and $c \geq 0$, any super-solution of (5) satisfies

$$\sup_{B_d} u^- \leq \sup_{\partial B_d} u^- + Cd \left(\left(\int_{B_d \cap \{u + M_\partial = \Gamma(u)\}} (f_0^+)^n \right)^{1/n} \right)^{\frac{1}{1+\alpha}} \tag{27}$$

where $M_\partial = \sup_{\partial B_d} u^-$, $\Gamma(u)$ is the convex hull of $\min(u + M_\partial, 0)$ extended by 0 on B_{2d} , and C is a constant (only) depending on $\|b\|_{L^n(B_d)}$, n , α and $\|c\|_\infty$.

Dávila, Felmer and Quaas pointed out to us that it can be obtained from ours. See below.

The Harnack inequality obtained in [3] is the following one

Theorem 7. (See [3, Theorems 3.1 and 3.2].) Under Assumption (H) and $c \geq 0$, any non-negative solution of (5) satisfies

$$\sup_B u \leq C \left(\inf_B u + \|f_0\|_{L^n(B)}^{\frac{1}{1+\alpha}} \right) \tag{28}$$

where C is a constant (only) depending on n , q , λ_F , Λ_F , $\|c\|_\infty$, α and $\|b\|_{L^q(Q_1)}$.

Remark 10. This result is proved in [3] only in dimension 2. Moreover, it is slightly more precise since it depends on q and $\|b\|_{L^q(Q_1)}$ instead of $\|b\|_{L^\infty(Q_1)}$.

Their results are not included in ours but they can be derived with little additional work. We mention that Birindelli and Demengel do not prove this Harnack inequality by proving first an ABP estimate.

Proof of Theorem 6. Dávila, Felmer and Quaas kindly explained to us the link between their result and our result. We slightly adapt their argument to get the general case.

Assumption (H2) implies $|p|^\alpha \mathcal{M}^-(X) \leq F_0(p, X) \leq |p|^\alpha \mathcal{M}^+(X)$.

- If $\alpha \geq 0$, (7) and (8) are satisfied for any $M_F > 0$ with $\sigma = |b|$, $f = \frac{f_0^+}{M_F^\alpha}$ and $\gamma_F u$ is replaced with $cu|u|^\alpha$. Moreover, (6) is satisfied for any $M_F > 0$ and with $\sigma = |b|$ and $g(x, u) = \frac{(f_0(x) + cu|u|^\alpha)^+}{M_F^\alpha}$. In particular, $g(x, -M_\partial) \leq \frac{f_0(x)^+}{M_F^\alpha}$ since $c \geq 0$. Hence, our result gives

$$\sup_{B_d} u^- \leq \sup_{\partial B_d} u^- + Cd \left(M_F + \frac{1}{M_F^\alpha} \left(\int_{B_d \cap \{u + M_\partial = \Gamma(u)\}} (f_0^+)^n \right)^{1/n} \right).$$

Optimizing with respect to $M_F > 0$ gives (27).

- If $\alpha = -\beta < 0$, then $F(x, u, p, X) \geq 0$ and $u \leq 0$ implies

$$\mathcal{M}^+(X) + |b(x)||p| + (f_0)_+ |p|^\beta \geq 0.$$

Now using

$$g(p) = |p|^{-\beta n} \left(|p|^{\frac{n(n-\beta)}{n-1}} + \mu^{\frac{n}{n-1}} \right)^{-n}$$

in the proof of Theorem 1 permits to conclude after very similar computations. \square

The Harnack inequality of [3] when $c = 0$ can be easily obtained from ours in any dimension (but not when $c \neq 0$). The case $c \neq 0$ can also be treated but it requires to slightly modify proofs.

7. Proofs

7.1. Proof of the weak Harnack inequality

Proof of the weak Harnack inequality (Theorem 2). The proof of the weak Harnack inequality is performed in four steps. First, the problem is reduced to the case of a cube Q of universal side-length (Lemma 6), then it is proved that non-negative super-solutions can be bounded from above on Q by a universal constant on a set of universal positive measure (Lemma 7). Next, the measures of all level sets of super-solutions (restricted to Q) are (universally) estimated from above. Finally, we prove the weak Harnack inequality in Q .

Step 1. As explained above, we first reduce the problem to a simpler one.

Lemma 6 (Reduction of the problem). Consider a non-negative super-solution u of (4) in Q_{2R} . Then there exist universal constants p_0, ε_0 and C satisfying

$$\left. \begin{aligned} \inf_{Q_{3r}} u &\leq 1 \\ \max(M_F, \gamma_F, \|f\|_{L^n(Q_R)}, \|\sigma\|_{L^q(Q_R)}) &\leq \varepsilon_0 \end{aligned} \right\} \Rightarrow \|u\|_{L^{p_0}(Q_r)} \leq C. \quad (29)$$

We now explain how to derive the weak Harnack inequality from it. Let v be a super-solution of (4) in $Q_{R/t}$ for some $t > 0$. We then define a function $v_s(y) = \frac{v(ty)}{V}$ with $V > 0$ and $t \in (0, 1)$ to be chosen later. Thanks to Lemma 2 with $x_0 = 0$, $M_0 = V$, $R_0 = R/t$, the new function v_s satisfies $F_s \geq 0$ in Q_R for a non-linearity F_s satisfying (A) and (7) with

$$M_s = \frac{tM_F}{V}, \quad \gamma_s = \gamma_F t^2, \quad \sigma_s(y) = t\sigma(ty), \quad f_s(y) = \frac{f^+(ty)}{V}.$$

Hence, if one chooses

$$V = \inf_{Q_{3r/t}} v + \delta + \varepsilon_0^{-1} \max(M_F, \|f\|_{L^n(Q_{R/t})}),$$

$$t = \left(\left(\frac{\|\sigma\|_{L^q(Q_{R/t})}}{\varepsilon_0} \right)^{q/(q-n)} + \left(\frac{\gamma_F}{\varepsilon_0} \right)^{1/2} + 1 \right)^{-1}$$

we obtain that v satisfies

$$\inf_{Q_{3r}} v_s \leq 1,$$

$$\max(M_s, \gamma_s, \|f_s\|_{L^n(Q_R)}, \|\sigma_s\|_{L^q(Q_R)}) \leq \varepsilon_0.$$

We thus can apply Lemma 6 and we obtain from (29) the following estimate (after letting $\delta \rightarrow 0$)

$$\|v\|_{L^{p_0}(Q_{r/t})} \leq C \left(\inf_{Q_{3r/t}} v + \max(M_F, \|f\|_{L^n(Q_{R/t})}) \right). \tag{30}$$

A standard covering procedure permits to get (19).

Step 2. In this step, we obtain a (universal) upper bound M for non-negative super-solutions in Q_R on a set of (universal) positive measure μ if the L^n -norm of f on Q_R , the L^q -norm of σ on Q_R , M_F and γ_F are (universally) small.

Lemma 7 (Upper bound on a subset of positive measure). *There exist universal constants $r, R > 0$, $\varepsilon_0 > 0$, $\mu \in (0, 1)$ and $M_B > 0$ such that for any non-negative super-solution u of (4) in Q_R , we have*

$$\left. \begin{array}{l} \inf_{Q_{3r}} u \leq 1 \\ \max(M_F, \gamma_F, \|f\|_{L^n(Q_R)}, \|\sigma\|_{L^q(Q_R)}) \leq \varepsilon_0 \end{array} \right\} \Rightarrow |\{u \leq M_B\} \cap Q_r| \geq \mu |Q_r|.$$

The proof of this lemma relies on the barrier function φ that we constructed in the preliminary section and on the ABP estimate applied to $w = u + \varphi$.

Proof of Lemma 7. Given $\varepsilon_0 > 0$ to be fixed later, we consider φ from Lemma 1 and define $w = u + \varphi$. We want to apply the ABP estimate (Theorem 1) to the function w on the ball B_R .

- First, $u \geq 0$ and $\varphi \geq 0$ on ∂B_R hence $M_\partial = \sup_{\partial B_R} w^- = 0$.
- Since $\inf_{Q_{3r}} u \leq 1$ and $\varphi \leq -2$ in Q_{3r} , we conclude that $\inf_{Q_{3r}} w \leq -1$; in other words, we have $\sup_{Q_{3r}} w^- \geq 1$.
- We also claim that w is a super-solution of an appropriate equation. More precisely, we claim that w satisfies (18) in $\{w \leq 0\} \cap B_R$ for some appropriate continuous functions \bar{f} and $\bar{\sigma}$.

Let us justify the last assertion and make precise what \bar{f} and $\bar{\sigma}$ are. We write

$$\begin{aligned} 0 &\leq F(x, u, Du, D^2u) \\ &= F(x, w - \varphi, Dw - D\varphi, D^2w - D^2\varphi) \\ &\leq F(x, w + M_B, Dw - D\varphi, D^2w - D^2\varphi). \end{aligned}$$

Assume next that $|Dw| \geq M_F + \varepsilon_0 =: \bar{M}_F$, $D^2w \geq 0$ (in the viscosity sense) and $w \leq 0$. Then $|Dw - D\varphi| \geq M_F$ and we obtain from (7) the following inequality

$$0 \leq \mathcal{M}^+(D^2w) - \mathcal{M}^-(D^2\varphi) + \sigma|Dw| + \gamma_F M_B + \sigma\varepsilon_0 + f$$

(we used the fact that \mathcal{M}^+ is subadditive and the relation between the two Pucci operators). Use next that $D^2w \geq 0$ and φ satisfies (15)

$$\lambda_F \Delta w \leq \sigma|Dw| + C_B \xi + \gamma_F M_B + \sigma\varepsilon_0 + f.$$

Hence (18) holds true with

$$\bar{\sigma} = \sigma \quad \text{and} \quad \bar{f}(x) = C_B \xi + \gamma_F M_B + \sigma\varepsilon_0 + f.$$

By using the ABP estimate for w and the properties listed above satisfied by this function, we obtain

$$1 \leq \sup_{B_R} w^- \leq 3e^{C_{ABP}(1 + \|\bar{\sigma}\|_{L^n(B_R)}^n)} R \left(\bar{M}_F + \left(\int_{\{\Gamma(w)=w\} \cap B_R} (\bar{f}^+)^n \right)^{1/n} \right)$$

where $\Gamma(w)$ is the convex hull of $\min(w, 0)$ after extending w to B_{2R} by setting $w \equiv 0$ outside B_R . We now use the fact that

$$\max(M_F, \gamma_F, \|f\|_{L^n(Q_R)}, \|\sigma\|_{L^q(Q_R)}) \leq \varepsilon_0,$$

together with definitions of \bar{f} , \bar{M}_F and the fact that $\text{supp } \xi \subset Q_r$ in order to get

$$1 \leq 3e^{C_{ABP}(1 + (\omega_n R)^{n(1-\frac{n}{q})} \varepsilon_0^n)} R (3\varepsilon_0 + (\omega_n R)^{1-\frac{n}{q}} \varepsilon_0^2 + \varepsilon_0 M_B + C_B |\{\Gamma(w) = w\} \cap Q_r|).$$

It is now enough to remark that

$$\{\Gamma(w) = w\} \subset \{w \leq 0\} \subset \{u \leq M_B\}$$

and to choose $\varepsilon_0 \in (0, 1)$ such that

$$3e^{C_{ABP}(1 + (\omega_n R)^{n(1-\frac{n}{q})} \varepsilon_0^n)} R (3\varepsilon_0 + (\omega_n R)^{1-\frac{n}{q}} \varepsilon_0^2 + \varepsilon_0 M_B) \leq \frac{1}{2}$$

to conclude. We used here that M_B is universal; in particular, it does not depend on ε_0 . \square

Step 3. We derive from the previous lemma (Lemma 7) an estimate of any level set of super-solutions u under consideration. Precisely, we use Lemma 2 together with the Calderón-Zygmund cube decomposition lemma (see Lemma 15 in Appendix A) in order to get the following result.

Lemma 8 (Estimate of the measure of level sets in Q_r). Let u be as in Lemma 7. Then there exist universal constants $\varepsilon > 0$ and $d > 0$ such that for all $t > 0$,

$$|\{u \geq t\} \cap Q_r| \leq dt^{-\varepsilon}. \tag{31}$$

The proof of Lemma 4.6 in [6] can be easily adapted (with minor changes). For the reader's convenience, a detailed proof is given in Appendix A.

Step 4. We finally explain how to derive Lemma 6. We first recall the following useful fact: if u is a non-negative function, then

$$\int_{Q_r} u^{p_0} = p_0 \int_0^{+\infty} t^{p_0-1} |\{u \geq t\} \cap Q_r| dt.$$

We can use the results of Lemmata 7 and 8: we thus choose $p_0 = \varepsilon/2$ where ε appears in (31) in order to get

$$\frac{1}{p_0} \int_{Q_r} u^{p_0} \leq \int_0^1 t^{\varepsilon/2-1} |Q_r| dt + \int_1^{+\infty} t^{\varepsilon/2-1} t^{-\varepsilon} dt =: C.$$

This achieves the proof of Lemma 6 and the proof of Theorem 2. \square

7.2. Proof of the local maximum principle

The proof of the local maximum principle is easily adapted from [6]. However, we give a detailed proof for the sake of completeness.

Proof of Theorem 3. The proof is divided in two steps. First, the problem is reduced to the case where the L^ε -norm of u is small; it is to be proven that u is bounded by a universal constant (Step 1). Then we explain how to get the universal bound (Steps 2 and 3).

Step 1. We state the lemma to be proven in Steps 2 and 3.

Lemma 9. Consider a sub-solution u of (4) in Q_R . Then there exists a universal constant $C > 0$ such that

$$\left. \begin{aligned} \|u^+\|_{L^\varepsilon(Q_r)} \leq d^{1/\varepsilon} \\ \max(M_F, \gamma_F, \|f\|_{L^q(Q_R)}, \|\sigma\|_{L^q(Q_R)}) \leq \varepsilon_0 \end{aligned} \right\} \Rightarrow \sup_{Q_{\frac{r}{4}}} u \leq C$$

where ε and d appear in Lemma 8.

We now explain how to derive Theorem 3 from this lemma. First, it is enough to get (20) for a particular p since the full result can be obtained by interpolation. In view of the previous lemma, we consider $p = \varepsilon$. By scaling u and by using a covering argument, we obtain the desired result.

Step 2. We remark that the assumption $\|u^+\|_{L^\varepsilon(Q_r)} \leq d^{1/\varepsilon}$ implies for all $t > 0$,

$$|\{u \geq t\} \cap Q_r| \leq t^{-\varepsilon} \int_{Q_r} (u^+)^{\varepsilon} \leq dt^{-\varepsilon}.$$

Remark that this estimate already appeared in the proof of the weak Harnack inequality; see (31) above. We next prove the following lemma.

Lemma 10. Consider a sub-solution u of (4) in Q_R satisfying (31) and F be such that

$$\max(M_F, \gamma_F, \|f\|_{L^n(Q_R)}, \|\sigma\|_{L^q(Q_R)}) \leq \varepsilon_0.$$

Then there exist universal constants $M_0 > 1$ and $\Sigma > 0$ such that

$$\left. \begin{array}{l} x_0 \in Q_{\frac{r}{2}}, \quad j \in \mathbb{N} \\ u(x_0) \geq \nu^{j-1} M_0 \end{array} \right\} \Rightarrow \sup_{Q_{l_j}(x_0)} u > \nu^j M_0$$

where $l_j = \Sigma \frac{M_0^{-\varepsilon/n}}{\nu^{\varepsilon j/n}} < \frac{r}{2}$ and $\nu = M_0/(M_0 - 1/2) > 1$.

Proof. We first choose Σ and M_0 such that

$$\Sigma M_0^{-\varepsilon/n} \leq \frac{r}{2}$$

so that $l_j < \frac{r}{2}$ and $Q_{l_j}(x_0) \subset Q_r$. We now argue by contradiction by assuming that $\sup_{Q_{l_j}(x_0)} u \leq \nu^j M_0$. We have to exhibit a contradiction.

On one hand, we have from (31) and the fact that $r < R$ and $l_j < r/2$

$$\left| \left\{ u \geq \nu^j \frac{M_0}{2} \right\} \cap Q_{\frac{l_j r}{R}}(x_0) \right| \leq d \nu^{-j\varepsilon} \left(\frac{M_0}{2} \right)^{-\varepsilon}. \tag{32}$$

On the other hand, since we have $\sup_{Q_{l_j}(x_0)} u \leq \nu^j M_0$ by assumption, we can consider the following transformation

$$T(y) = x_0 + \frac{l_j}{R} y$$

which defines a bijection between Q_R and $Q_{l_j}(x_0)$. The function ν defined on Q_R as follows

$$\nu(y) = \frac{\nu M_0 - \frac{u(T(y))}{\nu^{j-1}}}{(\nu - 1)M_0} \geq 0$$

thus satisfies $F_s(y, \nu, D\nu, D^2\nu) = 0$ in Q_R with F_s satisfying (A) and (7) with

$$\begin{aligned} M_s &= \frac{t}{\nu^{j-1}(\nu - 1)M_0} M_F, & \sigma_s(y) &= t\sigma(x_0 + ty), \\ \gamma_s &= t^2 \gamma_F, & f_s(y) &= \frac{t}{\nu^{j-1}(\nu - 1)M_0} t f(x_0 + ty) \end{aligned}$$

where $t = \frac{l_j}{R} < \frac{1}{2}$. It is clear that $\gamma_s \leq \gamma_F \leq \varepsilon_0$. Notice that

$$(\nu - 1)M_0 = \frac{M_0}{2M_0 - 1} > \frac{1}{2} > t$$

hence $M_S \leq M_F \leq \varepsilon_0$ and $f_S(y) \leq tf(x_0 + ty)$. We also have

$$\begin{aligned} \|\sigma_S\|_{L^q(Q_R)} &\leq t^{1-\frac{n}{q}} \|\sigma\|_{L^q(Q_R)} \leq \varepsilon_0, \\ \|f_S\|_{L^n(Q_R)} &\leq \|f\|_{L^n(Q_R)} \leq \varepsilon_0. \end{aligned}$$

Moreover, $v(0) = \frac{\nu M_0 - \frac{u(x_0)}{\nu^{j-1}}}{(\nu-1)M_0} \leq 1$ by assumption on u ; thus $\inf_{Q_{3r}} v \leq 1$. Hence, v satisfies the assumptions of Lemma 7 and we therefore obtain from Lemma 8 the following estimate

$$|\{v \geq M_0\} \cap Q_r| \leq dM_0^{-\varepsilon}.$$

We thus obtain

$$\left| \left\{ u \leq \nu^j \frac{M_0}{2} \right\} \cap Q_{\frac{l_j r}{R}}(x_0) \right| \leq \left(\frac{l_j}{R} \right)^n dM_0^{-\varepsilon}. \tag{33}$$

Combining (32) and (33), we thus obtain

$$\left(\frac{l_j r}{R} \right)^n \leq d\nu^{-j\varepsilon} \left(\frac{M_0}{2} \right)^{-\varepsilon} + \left(\frac{l_j}{R} \right)^n dM_0^{-\varepsilon}.$$

We also choose M_0 such that $dM_0^{-\varepsilon} \leq \frac{r^n}{2}$, and we obtain

$$\frac{1}{2} \left(\frac{l_j r}{R} \right)^n \leq d\nu^{-j\varepsilon} \left(\frac{M_0}{2} \right)^{-\varepsilon}.$$

Use now the definition of l_j and get

$$\frac{1}{2} \left(\frac{\Sigma r}{R} \right)^n \leq d2^\varepsilon.$$

We next choose $\Sigma > d^{\frac{1}{n}} 2^{\frac{\varepsilon+1}{n}} \frac{R}{r}$ in order to get a contradiction. \square

Step 3. We prove Lemma 9. By Step 2, we know that the sub-solution u satisfies the conclusion of Lemma 10. In particular, the series $\sum_j l_j$ converges and we can find a universal integer $j_0 \geq 1$ such that $\sum_{j \geq j_0} l_j \leq \frac{r}{4}$.

We now claim that $\sup_{Q_{\frac{r}{4}}} u \leq \nu^{j_0-1} M_0$. We argue by contradiction by assuming that this is not true and by exhibiting a contradiction. Let us assume that there exists $x_{j_0} \in Q_{\frac{r}{4}}$ such that $u(x_{j_0}) \geq \nu^{j_0-1} M_0$. Hence, we can apply Lemma 10 and we get a point x_{j_0+1} such that $|x_{j_0+1} - x_{j_0}|_\infty \leq l_{j_0}/2$ and $u(x_{j_0+1}) \geq \nu^{j_0} M_0$. By induction, we construct a sequence $(x_j)_{j \geq j_0}$ such that $|x_{j+1} - x_j| \leq l_j/2$ and $u(x_{j+1}) \geq \nu^j M_0$ as long as $x_j \in Q_{\frac{r}{2}}$. This is always the case since

$$|x_j|_\infty \leq |x_{j_0}|_\infty + \sum_{k=j_0}^{j-1} |x_{k+1} - x_k| \leq \frac{r}{8} + \frac{r}{8} \leq \frac{r}{4}.$$

We now get a contradiction since u is upper semi-continuous; indeed, it is bounded from above in $Q_{\frac{r}{2}}$ so it cannot satisfy $u(x_{j+1}) \geq \nu^j M_0$ for all $j \geq j_0$. The proof is now complete. \square

7.3. Proofs of Theorems 4 and 5

Proofs of Theorems 4 and 5. Both proofs rely on a transform of Cole–Hopf type in order to remove quadratic terms.

In order to understand why the exponential change of variables is the right one, we consider $v = h^{-1}(u)$ for some increasing convex function h and we remark that v satisfies

$$\mathcal{M}^+(D^2v) + \sigma(x)|Dv| + \frac{f^+(x)}{h'(v)} \geq 0$$

as soon as h satisfies $\lambda_F h'' - \sigma_2(h')^2 = 0$. We thus choose

$$h(t) = \frac{\lambda_F}{\sigma_2} \ln \left(1 - \frac{\sigma_2 t}{\lambda_F} \right)^{-1}.$$

We thus derive (23) from (19) by remarking that

$$\frac{1 - e^{-\frac{\sigma_2 \|u\|_{L^\infty(Q_1)}}{\lambda_F}}}{\frac{\sigma_2 \|u\|_{L^\infty(Q_1)}}{\lambda_F}} u \leq v \leq u$$

and $\frac{1}{h'(t)} = 1 - \frac{\sigma_2 t}{\lambda_F} \leq 1$.

We proceed in the same way in order to prove Theorem 5. Remark that we can assume without loss of generality that the solution is non-negative. \square

Acknowledgments

We are very grateful to Delarue for bringing our attention to this problem and for the fruitful discussions we had together. We also would like to thank Capuzzo-Dolcetta, Dávila, Felmer and Quaas for sending us their preprints and for their interest in our work and useful comments. In particular, the important remarks sent to us by Dávila, Felmer and Quaas permit us to improve the first version of this paper and to improve the results of Birindelli and Demengel.

Appendix A. Additional proofs

A.1. Proofs of Lemmata 3 and 4

In this paragraph, we explain how to prove Lemmata 3 and 4 by adapting the techniques of [17].

We first recall useful facts from convex analysis. The first one deals with the convex hull U^{**} of a function U .

Proposition 1. *Let Ω be a bounded convex open set and $U : \overline{\Omega} \rightarrow \mathbb{R}$ be lower semi-continuous. For $x \in \overline{\Omega}$, consider $(p, A) \in D^{2,-}U^{**}(x)$. Then there exist $x_1, \dots, x_q \in \overline{\Omega}$, $q \leq n + 1$, $\lambda_1, \dots, \lambda_q \in (0, 1]$, $\sum_{i=1}^q \lambda_i = 1$ such that*

$$\begin{cases} x = \sum_{i=1}^q \lambda_i x_i, \\ U^{**}(x) = \sum_{i=1}^q \lambda_i U(x_i). \end{cases} \tag{34}$$

Moreover U^{**} is linear on the convex hull of $\{x_1, \dots, x_q\}$. In particular, $A \leq 0$ for a.e. $x \in \{U = U^{**}\}$.

We next recall a result from [17] (see also [1]) about the subset of the convex hull U^{**} of a function U .

Proposition 2. (See [17, Proposition 3].) Let Ω be a bounded convex open set and $U : \overline{\Omega} \rightarrow \mathbb{R}$ be lower semi-continuous. For $x \in \overline{\Omega}$, consider $(p, A) \in D^{2,-}U^{**}(x)$. Consider x_i and λ_i such that (34) hold true. Then for every $\varepsilon > 0$, there are $A_i \in \mathbb{S}^{N-1}$, $i = 1, \dots, q$, such that

$$\begin{cases} (p, A_i) \in \overline{D}^{2,-}U(x_i), \\ A_\varepsilon \leq \square_{i=1}^q (\lambda_i^{-1} A_i) \end{cases} \quad (35)$$

where \square denotes the parallel sum of matrices. We recall that

$$(A \square B)\xi \cdot \xi = \inf_{\zeta \in \mathbb{R}^n} \{ A(\xi - \zeta) \cdot (\xi - \zeta) + B\zeta \cdot \zeta \}.$$

We next recall a (necessary and) sufficient condition for a function to be semi-concave.

Lemma 11. (See [1, Lemma 1].) Consider a bounded convex open set Ω and $U : \Omega \rightarrow \mathbb{R}$ a lower semi-continuous function. Assume that there exists $C > 0$ such that for all $x \in \Omega$ and all $(p, A) \in D^{2,+}U(x)$, $A \leq CI$. Then $U - C|\cdot|^2/2$ is concave.

We finally recall a useful approximation lemma from [1].

Lemma 12. (See [1].) Consider a convex set Ω and a convex function $V : \Omega \rightarrow \mathbb{R}$. For all $(p, A) \in D^{2,-}V(x)$, there exist $(x_n)_n$ and $(p_n, A_n) \in D^{2,-}V(x_n)$ such that $x_n \rightarrow x$, $p_n \rightarrow p$, $A_n \geq 0$ and $A \leq A_n + \frac{1}{n}$.

We now turn to the proofs of the two lemmata.

Proofs of Lemmata 3 and 4. The function $v = u + M_\partial$ is a super-solution of

$$G(x, v, Dv, D^2v) = 0$$

with $G(x, r, p, X) = F(x, r + M_\partial, p, X)$. Then $\Gamma(u)$ is the convex hull of the function $\min(v, 0)$.

We first reduce the problem to the study of subset of the function $\Gamma(u)$.

Lemma 13. Assume that $\Gamma(u)$ satisfies the following properties

$$\exists C > 0 / \forall x \in \mathcal{B}, \forall (p, A) \in D^{2,-}\Gamma(u)(x), \quad A \leq CI, \quad (36)$$

$$\begin{cases} \forall x \in \mathcal{B} \cap \{ \Gamma(u) = u + M_\partial \}, \forall (p, A) \in D^{2,-}\Gamma(u)(x), \\ A \leq \lambda_F^{-1}(\sigma(x)|p| + f^+(x))I, \end{cases} \quad (37)$$

$$\Gamma(u) \text{ is linear on } \mathcal{B} \setminus \{ x \in B_d : \Gamma(u) = u + M_\partial \}. \quad (38)$$

Then $\Gamma(u)$ satisfies conclusions of Lemmata 3 and 4.

Proof. Thanks to Lemma 11, Eq. (36) implies that $\Gamma(u)$ is semi-concave in \mathcal{B} . Since $\Gamma(u)$ is convex, this implies that $\Gamma(u)$ is $C^{1,1}$ in \mathcal{B} . Hence Lemma 3 is proved. We next remark that (38) implies Point 1 in Lemma 4. Eventually, (37) together with Alexandroff theorem permits to get Point 2. We recall that Alexandroff theorem asserts that a convex function is almost every twice differentiable. Hence the proof of Lemma 13 is now complete. \square

We now prove the following lemma in order to achieve the proof of Lemmata 3 and 4.

Lemma 14. *The function $\Gamma(u)$ satisfies (36), (37) and (38).*

Proof. We first remark that (38) is a consequence of Proposition 1 and of Alexandroff theorem.

We now turn to the proof of (36) and (37). Consider next $x \in \mathcal{B}$ and $(p, A) \in D^{2,-} \Gamma(u)(x)$. Notice that we cannot just prove (36) for a.e. $x \in \mathcal{B}$. In view of the definition of \mathcal{B} (see also Remark 4), we know that $|p| \geq M_F$. Thanks to Lemma 12, we can assume without loss of generality that $A \geq 0$. We now distinguish two cases.

Case 1: $x \in \mathcal{B} \cap \{\Gamma(u) = u + M_\partial\}$. In such a case, $(p, A) \in D^{2,-} \Gamma(u)(x) \subset D^{2,-} u(x)$, and since $|p| \geq M_F$, we have $F(x, u(x), p, A) \geq 0$. Now (6) yields

$$-\lambda_F \operatorname{Tr} A + \sigma(x)|p| + f^+(x) \geq 0$$

and since $A \geq 0$, we conclude that (37) holds true and the right-hand side is bounded in B_d since $\Gamma(u)$ is Lipschitz continuous and σ and f^+ are continuous.

Remark that the previous inequality also holds true for A such that $(p, A) \in \bar{D}^{2,-} \Gamma(u)(x)$, $A \geq 0$, since the equation is also satisfied for limiting semi-jets.

Case 2: $x \in \mathcal{B} \setminus \{\Gamma(u) = u + M_\partial\}$. There then exist $x_i \in \bar{B}_d$ and $\lambda_i \in (0, 1]$, $i = 1, \dots, q$, such that (34) holds true (where $U = u + M_\partial$). We know that there is at most one point x_i on ∂B_{2d} and the others are in B_d ; if not, $\Gamma(u) \equiv 0$ and there is nothing to prove. Moreover, $x_i \in \mathcal{B}$ for $i = 1, \dots, q$.

By Proposition 2, for any $\varepsilon > 0$, there exist q matrices $\lambda_i^{-1} A_i \geq A_\varepsilon \geq 0$ such that $\square_{i=1}^q \lambda_i^{-1} A_i \geq A_\varepsilon$ and $(p, A_i) \in \bar{D}^{2,-} u(x_i)$.

If there are no points on ∂B_{2d} , we deduce from Case 1 that for all i , $A_i \leq CI$ and $A_\varepsilon \leq CI$ follows.

If $x_q \in \partial B_{2d}$, say, then we deduce from (34) that $\lambda_q \leq 2/3$; hence, there exists $i \in \{1, \dots, q-1\}$ such that $\lambda_i \geq 1/3n$. For instance $i = 1$. Then we conclude that

$$A_\varepsilon \leq \frac{1}{\lambda_1} A_1 \leq 3nCI.$$

Passing to the limit on ε , we obtain $A \leq CI$ (for some new constant C). \square

A.2. Proof of Lemma 8

In order to prove Lemma 8, we need the Calderón–Zygmund cube decomposition such as stated in [6]. We thus first recall it. We use notation from [6]. Given $r > 0$, the cube Q_r is split in 2^n cubes of half side-length. We do the same with all the new cubes and we iterate the process. The cubes obtained in this way are called *dyadic cubes*. If Q is a dyadic cube of Q_r , \tilde{Q} denotes a dyadic cube such that Q is one of 2^n cubes obtained from \tilde{Q} . It is referred to as the predecessor of Q .

Lemma 15 (Cube decomposition). *Consider $r > 0$ and two measurable subsets $A \subset B \subset Q_r$. Consider $\delta \in (0, 1)$ such that*

- $|A| \leq \delta |Q_r|$;
- if Q is a dyadic cube of Q_r such that $|A \cap Q| > \delta |Q|$, then $\tilde{Q} \subset B$.

Then $|A| \leq \delta |B|$.

As far as the proof of this lemma is concerned, the reader is referred to [6]. We now turn to the proof of Lemma 8.

Proof of Lemma 8. We are going to prove the following estimate

$$|\{u \geq (M_B)^k\} \cap Q_r| \leq (1 - \mu)^k |Q_r| \tag{39}$$

where M_B and μ are given by Lemma 7. The reader can check that (31) derives from (39) with $d = (1 - \mu)^{-1}$ and $\varepsilon = -\ln(1 - \mu)/\ln M_B$.

We prove (39) by induction. Lemma 7 implies that (39) holds for $k = 1$. We now consider $k \geq 2$, we assume that (39) holds for $k - 1$ and we prove it for k . To do so, we are going to apply Lemma 15 with the two following sets $A \subset B \subset Q_r$

$$A = \{u > (M_B)^k\} \cap Q_r,$$

$$B = \{u > (M_B)^{k-1}\} \cap Q_r$$

and with $\delta = 1 - \mu$. Remark that $A \subset \{u > M_B\} \cap Q_r$; hence $|A| \leq (1 - \mu)|Q_r|$. It thus remains to prove that if Q is a dyadic cube of Q_r such that

$$|A \cap Q| > (1 - \mu)|Q| \tag{40}$$

then the predecessor \tilde{Q} of Q satisfies $\tilde{Q} \subset B$. Consider such a dyadic cube $Q = Q_{\frac{r}{2^i}}(x_0)$ and suppose that \tilde{Q} is not contained in B . Then there exists $\tilde{x} \in \tilde{Q}$ such that $u(\tilde{x}) \leq (M_B)^{k-1}$. We now use Lemma 2 with $R_0 = R$, $t_0 = \frac{1}{2^i}$ and $M_0 = (M_B)^{k-1}$ to get a rescaled function u_s satisfying $F_s = 0$ with F_s such that (7) holds with constants $M_s \leq M_F$, $\gamma_s \leq \gamma_F$ and functions f_s, σ_s satisfying $\|f_s\|_{L^n(Q_R)} \leq \|f\|_{L^n(Q_R)}$ and $\|\sigma_s\|_{L^q(Q_R)} \leq \|\sigma\|_{L^q(Q_R)}$. We thus can apply Lemma 7 if $\inf_{Q_{3r}} u_s \leq 1$. This is indeed the case

$$\inf_{Q_{3r}} u_s \leq \frac{u(\tilde{x})}{(M_B)^{k-1}} \leq 1.$$

Hence, $|Q \setminus A| > (1 - \mu)|Q|$ which contradicts (40). \square

A.3. Proof of Corollary 2

Proof of Corollary 2. We use the notation of [6]: for all $r \in (0, 1)$, $m_r = \inf_{Q_r} u$, $M_r = \sup_{Q_r} u$, $o_r = M_r - m_r = \text{osc}_{Q_r} u$. The non-negative functions $u - m_1$ and $M_1 - u$ satisfy equations $F_- = 0$, $F^+ = 0$ respectively for some non-linearities F_- and F^+ satisfying (7), (8) with f replaced with $f + \gamma_F M_1$. Hence, we can apply the Harnack inequality to $M_1 - u$ and $u - m_1$ and get

$$M_{1/2} - m_1 \leq C(m_{1/2} - m_1 + \max(M_F, \|f\|_{L^n(Q_1)} + \gamma_F |m_1|)),$$

$$M_1 - m_{1/2} \leq C(M_1 - M_{1/2} + \max(M_F, \|f\|_{L^n(Q_1)} + \gamma_F |M_1|))$$

where we can assume without loss of generality that $C > 1$. Adding these two inequalities and rearranging terms, we obtain

$$\text{osc}_{Q_{1/2}} u \leq \frac{C - 1}{C + 1} \text{osc}_{Q_1} u + 2 \max(M_F, \|f\|_{L^n(Q_1)} + \gamma_F \|u\|_{L^\infty(Q_1)}).$$

We now use Lemma 8.3 in [16] in order to get (22). \square

We shed light on the fact that Lemma 1 is a fortiori true for $M_B > 1$. This is important when proving Lemmata 7 and 8.

References

- [1] O. Alvarez, J.-M. Lasry, P.-L. Lions, Convex viscosity solutions and state constraints, *J. Math. Pures Appl.* (9) 76 (1997) 265–288.
- [2] I. Birindelli, F. Demengel, Comparison principle and Liouville type results for singular fully nonlinear operators, *Ann. Fac. Sci. Toulouse Math.* 13 (2004) 261–287.
- [3] I. Birindelli, F. Demengel, Eigenfunctions for singular fully non linear equations in unbounded regular domain, arXiv: 0904.1720.
- [4] L. Caffarelli, M.G. Crandall, M. Kocan, A. Świąch, On viscosity solutions of fully nonlinear equations with measurable ingredients, *Comm. Pure Appl. Math.* 49 (1996) 365–397.
- [5] L.A. Caffarelli, Interior a priori estimates for solutions of fully nonlinear equations, *Ann. of Math.* (2) 130 (1989) 189–213.
- [6] L.A. Caffarelli, X. Cabré, *Fully Nonlinear Elliptic Equations*, Amer. Math. Soc. Colloq. Publ., vol. 43, American Mathematical Society, Providence, RI, 1995.
- [7] I. Capuzzo-Dolcetta, F. Leoni, A. Vitolo, The Alexandrov–Bakelman–Pucci weak maximum principle for fully nonlinear equations in unbounded domains, *Comm. Partial Differential Equations* 30 (2005) 1863–1881.
- [8] M.G. Crandall, H. Ishii, P.-L. Lions, User's guide to viscosity solutions of second order partial differential equations, *Bull. Amer. Math. Soc. (N.S.)* 27 (1992) 1–67.
- [9] G. Dávila, P. Felmer, A. Quaas, Harnack inequality for singular fully nonlinear operators and some existence results, *Calc. Var. Partial Differential Equations*, in press.
- [10] G. Dávila, P. Felmer, A. Quaas, Alexandroff–Bakelman–Pucci estimate for singular or degenerate fully nonlinear elliptic equations, *CRAS* 347 (19–20) (2009) 1165–1168.
- [11] E. De Giorgi, Sulla differenziabilità e l'analiticità delle estremali degli integrali multipli regolari, *Mem. Accad. Sci. Torino Cl. Sci. Fis. Mat. Natur.* (3) 3 (1957) 25–43.
- [12] F. Delarue, Krylov and Safonov estimates for degenerate quasilinear elliptic PDEs, *J. Differential Equations* 248 (4) (2010) 924–951.
- [13] H. Federer, *Geometric Measure Theory*, Grundlehren Math. Wiss., Band 153, Springer-Verlag, New York, 1969.
- [14] K. Fok, A nonlinear Fabes–Stroock result, *Comm. Partial Differential Equations* 23 (1998) 967–983.
- [15] P. Fok, Some maximum principles and continuity estimates for fully nonlinear elliptic equations of second order, PhD thesis, University of California, Santa Barbara, 1996.
- [16] D. Gilbarg, N.S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, second ed., Grundlehren Math. Wiss. (Fundamental Principles of Mathematical Sciences), vol. 224, Springer-Verlag, Berlin, 1983.
- [17] C. Imbert, Convexity of solutions and $C^{1,1}$ estimates for fully nonlinear elliptic equations, *J. Math. Pures Appl.* (9) 85 (2006) 791–807.
- [18] H. Ishii, P.-L. Lions, Viscosity solutions of fully nonlinear second-order elliptic partial differential equations, *J. Differential Equations* 83 (1990) 26–78.
- [19] S. Koike, A. Świąch, Maximum principle and existence of L^p -viscosity solutions for fully nonlinear uniformly elliptic equations with measurable and quadratic terms, *NoDEA Nonlinear Differential Equations Appl.* 11 (2004) 491–509.
- [20] S. Koike, A. Świąch, Maximum principle for fully nonlinear equations via the iterated comparison function method, *Math. Ann.* 339 (2007) 461–484.
- [21] S. Koike, T. Takahashi, Remarks on regularity of viscosity solutions for fully nonlinear uniformly elliptic PDEs with measurable ingredients, *Adv. Differential Equations* 7 (4) (2002) 493–512.
- [22] N.V. Krylov, M.V. Safonov, An estimate for the probability of a diffusion process hitting a set of positive measure, *Dokl. Akad. Nauk SSSR* 245 (1979) 18–20.
- [23] N.V. Krylov, M.V. Safonov, A property of the solutions of parabolic equations with measurable coefficients, *Izv. Akad. Nauk SSSR Ser. Mat.* 44 (1980) 161–175, 239.
- [24] O.A. Ladyženskaja, V.A. Solonnikov, N.N. Ural'ceva, *Linear and Quasilinear Equations of Parabolic Type*, translated from the Russian by S. Smith, Transl. Math. Monogr., vol. 23, American Mathematical Society, Providence, RI, 1967.
- [25] J. Nash, Continuity of solutions of parabolic and elliptic equations, *Amer. J. Math.* 80 (1958) 931–954.
- [26] A. Quaas, B. Sirakov, Principal eigenvalues and the Dirichlet problem for fully nonlinear elliptic operators, *Adv. Math.* 218 (2008) 105–135.
- [27] J. Serrin, Local behavior of solutions of quasi-linear equations, *Acta Math.* 111 (1964) 247–302.
- [28] N.S. Trudinger, Local estimates for subsolutions and supersolutions of general second order elliptic quasilinear equations, *Invent. Math.* 61 (1980) 67–79.
- [29] N.S. Trudinger, Comparison principles and pointwise estimates for viscosity solutions of nonlinear elliptic equations, *Rev. Mat. Iberoamericana* 4 (1988) 453–468.